

Motivic construction of cohomological invariants

NIKITA SEMENOV

Abstract

The norm varieties and the varieties with special correspondences play a major role in the proof of the Bloch-Kato Conjecture by M. Rost and V. Voevodsky. In the present paper we show that a variety which possesses a special correspondence is a norm variety. The proof heavily uses motivic cohomology and the category of motives of Voevodsky.

As an unexpected application we give a positive answer to a problem of J.-P. Serre about groups of type E_8 over $\mathbb{Q}$. Apart from this we include the proof of the Voevodsky conjecture about ν_n -varieties which is due to A. Vishik. This result plays an important role in our proofs.

1 Introduction

Let us recall some recent developments in the topics which are relevant for this paper.

1.1 Cohomological invariants

The study of cohomological invariants was initiated by J.-P. Serre in the 90'ies. Serre conjectured the existence of an invariant in $H^3(k, \mathbb{Q}/\mathbb{Z}(2))$ of G -torsors, where G is a simply-connected semisimple algebraic group over a field k . This invariant was constructed by M. Rost and is now called the *Rost invariant* of G (see [Inv, 31.B and pp. 448–449]).

Nowadays there exist numerous constructions and estimations of cohomological invariants for different classes of algebraic objects (see e.g. [GMS]). Nevertheless, the most constructions of cohomological invariants rely on a

specific construction of the object under consideration. Unfortunately, for many groups, like E_8 , there is no classification and no general construction so far.

1.2 Chow motives of projective homogeneous varieties

Another direction which we discuss now is the theory of motives of projective homogeneous varieties. Typical examples of such varieties are projective quadrics and Severi-Brauer varieties.

The study of projective homogeneous varieties was initiated by Rost when he provided a motivic decomposition of a Pfister quadric used later in the proof of the Milnor conjecture (see [Ro98]). The motives of Severi-Brauer varieties were studied by Karpenko and the motives of general quadrics by Izhboldin, Karpenko, Merkurjev, and Vishik.

A systematical theory of motives of general projective homogeneous varieties was developed in a series of our papers with a culmination in [PSZ] and [PS10] where the structure of the motive of a generically split homogeneous variety was established. Moreover, in some cases we provided an alternative construction of *generalized Rost motives* as indecomposable direct summands of generically split varieties (see [PSZ, §7]).

The technique developed in our papers does not use cohomological invariants of algebraic groups and of projective homogeneous varieties at all, and our arguments don't rely on the classification or on a construction of algebraic groups.

1.3 The Bloch-Kato Conjecture

The Bloch-Kato conjecture (more precisely its proof) provides a bridge between cohomological invariants and motives.

The Bloch-Kato conjecture says that for any n and any prime number p the norm residue homomorphism

$$K_n^M(k)/p \rightarrow H^n(k, \mu_p^{\otimes n})$$

$$\{a_1, \dots, a_n\} \mapsto (a_1) \cup \dots \cup (a_n)$$

between the Milnor K-theory and the Galois cohomology is an isomorphism.

The case $p = 2$ of the Bloch-Kato conjecture is known as the Milnor conjecture.

The proof of the Bloch-Kato conjecture goes as follows. Given a pure symbol u in $H^n(k, \mu_p^{\otimes n})$ M. Rost constructs a splitting variety X_u for u with some additional properties (If $p = 2$ one takes norm quadrics, for $n = 2$ one takes Severi-Brauer varieties, for $p = 3$ the construction is based on the Merkurjev-Suslin varieties (see [S08] for an explicit construction for $p = n = 3$), for $p \geq 5$ the construction is implicit). Then V. Voevodsky using the category of his motives and symmetric powers splits off a certain direct summand M_u from the motive of X_u (see [Vo03]). The motive M_u is called the *generalized Rost motive*.

Later Rost was able to simplify Voevodsky's construction and found the same motive M_u staying inside the category of classical Chow motives (see [Ro07]). His main idea was to produce an algebraic cycle on X_u called a *special correspondence* (see Definition 4.2), forget about the symbol u and use the special correspondence to construct the motive M_u . This motive is used then to construct exact triangles in the category of Voevodsky's motives. These triangles essentially involve mixed motives, in particular, deviate from the category of classical Chow motives, and are used (among other results of Rost and Voevodsky) to finish the proof of the Bloch-Kato conjecture.

The symbol u is a cohomological invariant of the variety X_u (see [Vo03, Theorem 7.3]). Apart from this, one can notice that the special correspondences of Rost resemble the algebraic cycles that we construct in the proof of our motivic decompositions (see [Ro07, Lemma 5.2] and [PSZ, Lemma 4.15]).

Summarizing: The technique developed by Rost and Voevodsky in the proof of the Bloch-Kato conjecture gives a way to produce algebraic cycles and motives out of cohomological invariants of varieties.

1.4 Conclusion

The discussion in the above subsections leads to the following conjecture. Namely, there should exist a way back from motives to cohomological invariants. Note that an evidence of this conjecture appears already in papers of O. Izhboldin and A. Vishik [IzhV00] and [Vi98], where the case of quadrics was treated.

In the present paper we show in a constructive way that the varieties which possess a special correspondence of Rost admit cohomological invariants.

This result has several unexpected applications in the theory of algebraic groups. In the present paper we discuss just one of them. Namely, we

construct a cohomological invariant in $H^5(k, \mathbb{Z}/2)$ for any group of type E_8 whose Rost invariant is trivial. In turn, this invariant allows to give a positive answer to a question of Serre about compact groups of type E_8 (see the last section). This finishes Serre's program on classification of finite subgroups in Chevalley groups (see [GS09]).

1.5 Structure of the paper

First we recall some results on algebraic cobordism of Levine-Morel and ν_n -varieties. We need them to prove the Voevodsky conjecture. The proof belongs to A. Vishik, and we represent it here with his kind permission. Then we recall basic properties of the category of motives of Voevodsky and the Rost construction of special correspondences. Finally, we provide some general results which allow to “compress” varieties and finish the proofs of all main results of the present paper.

2 Algebraic cobordism and ν_n -varieties

2.1 (Lazard ring). Fix a prime p and a field k with $\text{char } k = 0$. Consider the algebraic cobordism Ω^* of Levine-Morel (see [LM]) and the Lazard ring $\mathbb{L} = \Omega(\text{Spec } k)$. This is a graded ring generated by the cobordism classes of maps $[X] = [X \rightarrow \text{Spec } k] \in \mathbb{L}^{-\dim X}$, where X is a smooth projective variety over k . It is known that this ring is isomorphic to the polynomial ring with integer coefficients on a countable set of variables.

2.2 (Characteristic numbers). Given a partition $J = (l_1, \dots, l_r)$ of arbitrary length $r \geq 0$ with $l_1 \geq l_2 \geq \dots \geq l_r > 0$ one can associated with it a characteristic class $c_J(X) \in \text{CH}^{|J|}(X)$ ($|J| = \sum_{i \geq 1} l_i$) of X as follows: Let $P_J(x_1, \dots, x_r)$ be the smallest symmetric polynomial containing the monomial $x_1^{l_1} \dots x_r^{l_r}$. We can express P_J as a polynomial on the standard symmetric function $\sigma_1, \dots, \sigma_r$ as $P_J(x_1, \dots, x_r) = Q_J(\sigma_1, \dots, \sigma_r)$ for some polynomial Q_J . Let $c_i = c_i(-T_X)$ denote the i -th Chern class of the virtual normal bundle of X . Then

$$c_J(X) = Q_J(c_1, \dots, c_r).$$

The degrees of the characteristic classes are called the characteristic numbers.

If $J = (1, \dots, 1)$ (i times), $c_J(X) = c_i(-T_X)$ is the usual Chern class. We denote $b_i(X) := c_{(p-1, \dots, p-1)}(X)$ (i times). In particular, if $p = 2$ then

$b_i(X) = c_i(-T_X)$. If $J = (p^n - 1)$, we write $c_J(X) = s_{p^n-1}(X)$. The class $s_d(X)$ is called the d -th *Milnor class* of X . Fortunately, if $d := \dim X = p^n - 1$ we have

$$b_d(X) = \pm s_d(X) \bmod pI(X) + p^2\mathbb{Z}_{(p)},$$

where $I(X) = \deg(\mathrm{CH}_0(X)) \subset \mathbb{Z}$ (see [Ro07, Lemma 9.13]).

2.3 Definition (ν_n -varieties). Let p be a fixed prime.

1. A smooth projective variety X is called a ν_n -variety if $\dim X = p^n - 1$, all characteristic numbers of X are divisible by p , and

$$s_{\dim X}(X) \neq 0 \bmod p^2.$$

2. A smooth projective variety X is called a $\nu_{\leq n}$ -variety if X is a ν_n -variety and for all $0 \leq i \leq n$ there exists a ν_i -variety Y_i and a morphism $Y_i \rightarrow X$.

2.4 Example. For $p = 2$ a typical example of a ν_n -variety is a smooth projective $(2^n - 1)$ -dimensional quadric (see [Vo01, Proposition 3.4]).

2.5 (Landweber-Novikov operations). Further on one can construct the Landweber-Novikov operations on the cobordism ring $\Omega^*(V)$ for a smooth quasi-projective variety V . Given a partition J as above we write

$$S_{L-N}^J: \Omega^*(V) \rightarrow \Omega^{*+|J|}(V)$$

for the operation which maps the cobordism class $[f: X \rightarrow V] \in \Omega^*(V)$ to $f_*(Q_J(c_1^\Omega, \dots, c_r^\Omega))$ where $c_i^\Omega = c_i^\Omega(-T_X + f^*(T_V))$ are the cobordism Chern classes (see [Vi07, Definition 2.12]).

Since Ω^* is a universal cohomology theory, there exists a canonical map $\mathrm{pr}: \Omega^* \rightarrow \mathrm{CH}^*$ which turns out to be surjective. For a particular partition $J = (p-1, \dots, p-1)$ (i times) denote $S_{L-N}^i := S_{L-N}^J$. Then the following diagram commutes:

$$\begin{array}{ccccc} \Omega^*(V) & \xrightarrow{\mathrm{pr}} & \mathrm{CH}^*(V) & \longrightarrow & \mathrm{Ch}^*(V) \\ S_{L-N}^i \downarrow & & & & \downarrow S^i \\ \Omega^{*+(p-1)i}(V) & \xrightarrow{\mathrm{pr}} & \mathrm{CH}^{*+(p-1)i}(V) & \longrightarrow & \mathrm{Ch}^{*+(p-1)i}(V) \end{array}$$

where $\text{Ch}^* := \text{CH}^* / p$ and S^i denotes the i -th Steenrod operation in the Chow theory (see [Br03a]). In particular, since $S^i|_{\text{Ch}^m(V)} = 0$ for $i > m$, the image

$$\text{pr} \circ S_{L-N}^i(\Omega^m(V))$$

is divisible by p for such m . Moreover, the total operation $S_{L-N} := \sum_i S_{L-N}^i$ is a ring homomorphism (see [Za09], [Vi07], and [LM]).

For an integer n let $I(p, n)$ denote the ideal of $\mathbb{L}$ generated by the varieties of dimension $\leq p^n - 1$ with all characteristic numbers divisible by p . This is a prime ideal invariant under the action of the Landweber-Novikov operations on $\mathbb{L}$. By definition $I(p, n-1) \subset I(p, n)$. Moreover, as an $\mathbb{L}$ -ideal $I(p, n)$ is generated by $I(p, n-1)$ and by any ν_n -variety.

2.6 Lemma (Voevodsky's conjecture). *A ν_n -variety is a $\nu_{\leq n}$ -variety.*

Proof. (A. Vishik) It is well known that a smooth hypersurface of degree p in $\mathbb{P}^{p^n}$ is a ν_n -variety (see [Vo01, Proposition 3.4]). Denote its class in $\mathbb{L}$ as u . Moreover, a direct computation shows that $S_{L-N}^{p^{n-1}}(u)$ is a multiple of the class of a hypersurface of degree p in $\mathbb{P}^{p^{n-1}}$ with a coefficient not divisible by p . In particular, $S_{L-N}^{p^{n-1}}(u)$ is a ν_{n-1} -element.

Let now v_n be a ν_n -element in $\mathbb{L}$. Then $(av_n - u) \in I(p, n-1)$ for some $a \in \mathbb{Z}$ coprime to p . To conclude that v_n is a ν_{n-1} -element, it suffices to show that $S_{L-N}^{p^{n-1}}(y) \in I(p, n-2)$ for any $y \in I(p, n-1)$. We can write

$$y = v_{n-1}x_{n-1} + v_{n-2}x_{n-2} + \dots + v_0x_0$$

where v_i is a ν_i -element and $x_i \in \mathbb{L}$. Since the total operation S_{L-N} is multiplicative, we see that

$$S_{L-N}^{p^{n-1}}(v_{n-2}x_{n-2} + \dots + v_0x_0) \in I(p, n-2).$$

Moreover, $S_{L-N}^i(v_{n-1}) \in I(p, n-2)$ for all $i > 0$. Finally, we have the term

$$S_{L-N}^0(v_{n-1})S_{L-N}^{p^{n-1}}(x_{n-1}) = v_{n-1}S_{L-N}^{p^{n-1}}(x_{n-1}).$$

But $S_{L-N}^{p^{n-1}}(x_{n-1})$ is divisible by p , since this Landweber-Novikov operation induces the Steenrod operation $S^{p^{n-1}}$ on the Chow group. Thus,

$$S_{L-N}^{p^{n-1}}(y) \in I(p, n-2),$$

and we are done. □

3 Category DM

The base field k is assumed to have characteristic zero, and all varieties are assumed to be irreducible.

If $A \rightarrow B$ is a morphism in a triangulated category, then there exists a unique up to isomorphism object C with an exact triangle of the form $A \rightarrow B \rightarrow C$. We write $C = \text{Cone}(A \rightarrow B)$.

The following lemma is a standard property of triangulated categories.

3.1 Lemma. *Assume we are given an exact (or distinguished) triangle*

$$A \rightarrow B \rightarrow C \rightarrow A[1]$$

in a triangulated category. Then for any object U in this category the following induced sequences are exact:

$$\dots \rightarrow \text{Hom}(C, U) \rightarrow \text{Hom}(B, U) \rightarrow \text{Hom}(A, U) \rightarrow \text{Hom}(C[-1], U) \rightarrow \dots$$

$$\dots \rightarrow \text{Hom}(U, A) \rightarrow \text{Hom}(U, B) \rightarrow \text{Hom}(U, C) \rightarrow \text{Hom}(U, A[1]) \rightarrow \dots$$

3.2 Definition. Let X be a smooth projective variety over k . As $\check{C}(X)$ we denote the *standard simplicial scheme* associated with X :

$$X \rightrightarrows X \times X \rightrightarrows X \times X \times X \dots$$

In the present paper we work in the category $\text{DM} := \text{DM}_-^{eff}(k)$ of effective motives of V. Voevodsky over k with $\mathbb{Z}/p$ -coefficients, where p is a prime number. It is well-known that there are functors from the category of smooth schemes over k and from the category of smooth simplicial schemes over k to DM . Therefore we can speak about the motives of smooth schemes and smooth simplicial schemes.

The category of classical Grothendieck-Chow motives (with $\mathbb{Z}/p$ -coefficients) is a full subcategory of DM closed under direct summands. Its objects are called *pure motives*. For an integer b we set $\{b\} := (b)[2b]$. For example, if $M \in \text{DM}$, then $M\{b\}[1]$ means $M(b)[2b+1]$ and $M\{b\}$ is a *Tate twist* of M . For $M \in \text{DM}$ its *motivic cohomology*:

$$H^{d,c}(M, \mathbb{Z}/p) = \text{Hom}(M, \mathbb{Z}/p(c)[d]).$$

We recall now some well-known properties of the category DM which are due to Voevodsky.

3.3 Lemma ([Vo95a, Proposition 4.2.9] and [Vo95b, Proposition 2.7]). *Let X be a smooth projective variety over k . Then*

$$\mathrm{Hom}(X, \mathbb{Z}/p(i)[j]) = \mathrm{CH}^i(X, 2i - j) \otimes \mathbb{Z}/p$$

is the higher Chow group of X and

$$\mathrm{Hom}(X, \mathbb{Z}/p(i)[2i]) = \mathrm{CH}^i(X) \otimes \mathbb{Z}/p =: \mathrm{Ch}^i(X)$$

is the Chow group of X . If $X = \mathrm{Spec} k$, then additionally

$$\mathrm{Hom}(X, \mathbb{Z}/p(i)[i]) = K_i^M(k)/p$$

is the Milnor K -theory of $k \bmod p$.

3.4 Lemma ([Vi98, Theorem 2.3.4]). *Let X be a smooth projective variety over k and let $\mathcal{X}$ be the motive of $\check{C}(X)$. The natural morphism $\mathcal{X} \rightarrow \mathrm{Spec} k$ is an isomorphism if and only if X has a zero-cycle of degree coprime to p .*

3.5 Lemma ([Vo96, Corollary 2.3]). *Let X be a smooth projective variety over k and M a direct summand of the motive of X . Then*

$$\mathrm{Hom}(M, \mathbb{Z}/p(c)[d]) = 0$$

for $d - c > \dim X$ and for $d > 2c$.

3.6 Lemma (Cancellation; [Vo02, Corollary 4.10]). *Let $A, B \in \mathrm{DM}$. Then for all $i, j \geq 0$, we have*

$$\mathrm{Hom}(A, B) = \mathrm{Hom}(A(i)[j], B(i)[j]).$$

3.7 Definition. A localizing subcategory of DM generated by an object $A \in \mathrm{DM}$ is a minimal triangulated subcategory closed under direct summands, arbitrary direct sums, tensor products and Tate twists, and containing A .

3.8 Lemma ([Vo96, Lemma 3.8 and Lemma 4.7], [Vi98, Theorem 2.3.2]). *Let X be a smooth projective variety over k and M an object of the localizing subcategory of DM generated by X . Denote as $\mathcal{X}$ the motive of $\check{C}(X)$. Then*

1. *the natural morphism $M \otimes \mathcal{X} \rightarrow M$ is an isomorphism,*
2. *and the natural homomorphism $\mathrm{Hom}(M, \mathcal{X}\{i\}) \rightarrow \mathrm{Hom}(M, \mathbb{Z}/p\{i\})$ is an isomorphism for any $i \geq 0$.*

In DM there is an action of the *Milnor operations* for smooth simplicial schemes U

$$Q_i: \operatorname{Hom}(U, \mathbb{Z}/p(r)[s]) \rightarrow \operatorname{Hom}(U, \mathbb{Z}/p(r + p^i - 1)[s + 2p^i - 1]).$$

In the same way there is an action of the Milnor operations on the *reduced motivic cohomology* $\widetilde{H}^{*,*}(U, \mathbb{Z}/p)$ for smooth pointed simplicial schemes U . They satisfy the following properties:

3.9 Lemma ([Vo96, Theorem 3.17]). *Let U be a smooth simplicial scheme (resp. smooth pointed simplicial scheme) and $u, v \in H^{*,*}(U, \mathbb{Z}/p)$ (resp. $u, v \in \widetilde{H}^{*,*}(U, \mathbb{Z}/p)$). Then*

1. $Q_i^2 = 0$;
2. $Q_i Q_j + Q_j Q_i = 0$;
3. $Q_i(uv) = Q_i(u)v + uQ_i(v) + \sum \xi^{n_j} \varphi_j(u) \psi_j(v)$, where $n_j > 0$, φ_j and ψ_j are some homogeneous cohomological operations of bidegrees $(*, *)$ with $*' > 2* \geq 0$, and ξ is the class of -1 in $\operatorname{Hom}(\mathbb{Z}/p, \mathbb{Z}/p(1)[1]) = k^\times / k^{\times p}$. (In particular, $\xi = 0$ if p is odd).

Thus, the following sequence is a complex:

$$\widetilde{H}^{*-2p^i+1, *'-p^i+1}(U, \mathbb{Z}/p) \xrightarrow{Q_i} \widetilde{H}^{*,*'}(U, \mathbb{Z}/p) \xrightarrow{Q_i} \widetilde{H}^{*+2p^i-1, *'+p^i-1}(U, \mathbb{Z}/p). \quad (1)$$

3.10 Definition. The cohomology groups of complex (1) are called *Margolis motivic cohomology groups* and are denoted as $\widetilde{HM}_i^{*,*'}(U)$.

The following lemma is of particular importance for us.

3.11 Lemma. *Let X be a ν_n -variety over k and $\mathcal{X}$ be the motive of $\check{C}(X)$. Denote as $\widetilde{\mathcal{X}} = \operatorname{Cone}(\mathcal{X} \rightarrow \mathbb{Z}/p)$ the cone of the natural projection. Then for all $0 \leq i \leq n$ and all $*, *' \in \mathbb{Z}$*

$$\widetilde{HM}_i^{*,*'}(\widetilde{\mathcal{X}}) = 0.$$

Proof. The lemma immediately follows from [Vo03, Lemma 4.3] and Lemma 2.6. $\square$

3.12 Lemma ([Vo96, Corollary 2.2]). *We have $\operatorname{Hom}(\mathcal{X}, \mathbb{Z}/p(c)[d]) = 0$ if $c < 0$.*

3.13 Lemma ([IzhV00, Proof of Sublemma 6.2]). *Let X be a smooth projective variety and $\mu' \in \mathrm{Hom}(\mathcal{X}, \mathcal{X}\{b\}[1])$ for some b . Let $\mu \in \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p\{b\}[1])$ be the image of μ' under the natural map of Lemma 3.8(2).*

Then for all c, d the pull-back

$$(\mu')^*: \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p(c)[d]) \rightarrow \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p(c+b)[d+2b+1])$$

coincides with the multiplication by μ .

The next two lemmas follow from the Bloch-Kato Conjecture.

3.14 Lemma ([Vo03, Lemma 6.6]). *Let X be a smooth projective variety over k , let $\mathcal{X}$ be the motive of $\check{C}(X)$, and $\tilde{\mathcal{X}} = \mathrm{Cone}(\mathcal{X} \rightarrow \mathrm{Spec} k)$.*

Then

$$\tilde{H}^{d,c}(\tilde{\mathcal{X}}, \mathbb{Z}/p) = 0$$

for all $d \leq c + 1$.

3.15 Lemma ([Ro07, Lemma 2.1]). *Let X be a smooth projective (irreducible) variety over k , $n \geq 2$, and let $\mathcal{X}$ be the motive of $\check{C}(X)$. Then there is a natural exact sequence:*

$$0 \rightarrow H^{n,n-1}(\mathcal{X}, \mathbb{Z}/p) \rightarrow H_{et}^n(k, \mu_p^{\otimes(n-1)}) \rightarrow H_{et}^n(k(X), \mu_p^{\otimes(n-1)}).$$

In particular, any element of $H^{n,n-1}(\mathcal{X}, \mathbb{Z}/p)$ can be identified with an element of the Galois cohomology $H_{et}^n(k, \mu_p^{\otimes(n-1)})$.

Next we recall several facts about generically split motives and Rost nilpotency.

3.16 Definition. A motive M over k is called *generically split*, if there exists a smooth projective k -variety X and a number l such that M is a direct summand of the motive $X\{l\}$ and $M_{k(X)}$ is a direct sum of twisted Tate motives over $k(X)$.

3.17 Definition. Let M be a pure motive over k . We say that *Rost nilpotence* holds for M if for any field extension K/k the kernel of the natural map

$$\mathrm{End}_k(M) \rightarrow \mathrm{End}_K(M_K)$$

consists of nilpotent correspondences.

3.18 Lemma ([ViZ08, Proposition 3.1]; see also [Br03b, Theorem 3.1]). *Rost nilpotence holds for all generically split motives.*

4 Rost's construction

We assume that $\text{char } k = 0$ and k contains a primitive p -th root of unity ζ_p .

4.1 Definition (Norm varieties). Let $u \in H_{\text{et}}^n(k, \mu_p^{\otimes n})$. A smooth projective irreducible variety X over k is called a norm variety for u if X is a ν_{n-1} -variety and $u_{k(X)} = 0$.

If u is a pure symbol, i.e., $u = (a_1) \cup \dots \cup (a_n)$ for some $a_i \in H^1(k, \mu_p) \simeq k^\times/k^{\times p}$, then it was shown by M. Rost that a norm variety for u exists.

4.2 Definition (Rost). Let n be a positive integer, p a prime, $b = \frac{p^{n-1}-1}{p-1}$, and X a smooth projective irreducible variety over k of dimension $p^{n-1} - 1$. Consider the complex

$$\text{CH}^b(X) \xrightarrow{\pi_0^* - \pi_1^*} \text{CH}^b(X \times X) \xrightarrow{\pi_0^* - \pi_1^* + \pi_2^*} \text{CH}^b(X \times X \times X)$$

where π_i is the i -th projection in the diagram

$$X \xleftarrow{\quad} X \times X \xleftarrow{\quad} X \times X \times X.$$

Let $\rho \in \text{CH}^b(X \times X)$. Define

$$c(\rho) := (\pi_0)_*(\rho^{p-1}) \in \text{CH}^0(X) = \mathbb{Z}.$$

The cycle ρ is called a *special correspondence* of type (n, p) for X if

$$(\pi_0^* - \pi_1^* + \pi_2^*)(\rho) = 0$$

and

$$c(\rho) \neq 0 \pmod{p}.$$

M. Rost showed that a variety X which possesses a special correspondence and has no zero-cycles of degree coprime to p is a ν_{n-1} -variety (see [Ro07, Section 9]).

The following table gives some examples of norm varieties (cf. [Ro02]).

p	n	symbol	norm variety
2	any	$(a_1) \cup \dots \cup (a_n)$	the projective quadric given by $q = 0$ where $q = \langle\langle a_1, \dots, a_{n-1} \rangle\rangle \perp \langle -a_n \rangle$.

2	5	f_5	the variety of singular trace zero lines in a reduced Albert algebra with the cohomological invariant f_5 ; this variety is a twisted form of F_4/P_4 .
any	2	$(a) \cup (b)$	Severi-Brauer variety $\text{SB}(A)$ where $A = \langle x, y \mid x^p = a, y^p = b, xy = \zeta_p yx \rangle$.
3	3	g_3	the twisted form of a hyperplane section of $\text{Gr}(3, 6)$ defined in [S08]; this variety is closely related to Albert algebras with the Rost invariant g_3 .
5	3	h_3	the variety is related to groups of type E_8 with the Rost invariant h_3 ; it can be constructed from projective E_8 -homogeneous varieties using the algorithm of Section 7.
any	3	$(a) \cup (b) \cup (c)$	any smooth compactification of the Merkurjev-Suslin variety $\text{MS}(a, b, c) = \{\alpha \in A \mid \text{Nrd}_A(\alpha) = c\}$ where A is as above, and Nrd_A is its reduced norm.
3	4	$(a) \cup (b) \cup (c) \cup (d)$	any smooth compactification of the variety $\{\alpha \in J \mid N_J(\alpha) = d\}$ where J is the Albert algebra obtained from the 1st Tits construction out of A and c with A as above, and N_J is the cubic form on J .

We recall now the construction of the special correspondences of Rost out of a symbol. Let n be an integer, p a prime, $u \in H^n(k, \mu_p^{\otimes(n-1)})$ an element in the Galois cohomology of the field k , and X a smooth projective variety such that $u_{k(X)} = 0$.

By Lemma 3.15 the element u can be identified with an element $\delta \in H^{n, n-1}(\mathcal{X}, \mathbb{Z}/p)$ of the motivic cohomology of the standard simplicial scheme $\mathcal{X}$ associated with X .

Define now $\mu = \tilde{Q}_0 \circ Q_1 \circ \dots \circ Q_{n-2}(\delta) \in H^{2b+1, b}(\mathcal{X}, \mathbb{Z})$, where Q_i are the Milnor operations, $\tilde{Q}_0$ is the integral valued Bockstein, and $b = \frac{p^{n-1}-1}{p-1}$.

Consider now the *skeleton filtration* on $\mathcal{X}$. Let $Y = \text{Cone}(X \rightarrow \mathcal{X})$ be the cone of the natural map. Then there is another natural map $(X \times X)[1] \rightarrow Y$ such that the composition $(X \times X)[1] \rightarrow Y \rightarrow X[1]$ equals $\pi_0 - \pi_1$.

Let $Z = \text{Cone}((X \times X)[1] \rightarrow Y)$. Then again there is a natural map

$(X \times X \times X)[2] \rightarrow Z$ such that the composition $(X \times X \times X)[2] \rightarrow Z \rightarrow (X \times X)[2]$ equals $\pi_0 - \pi_1 + \pi_2$.

By Lemma 3.1 these exact triangles give the long exact sequences for motivic cohomology with integral coefficients:

$$H^{2b,b}(X) \rightarrow H^{2b+1,b}(Y) \rightarrow H^{2b+1,b}(\mathcal{X}) \rightarrow H^{2b+1,b}(X)$$

and

$$H^{2b+1,b}(Y) \rightarrow H^{2b,b}(X \times X) \xrightarrow{\alpha} H^{2b+2,b}(Z).$$

Note now that $H^{2b,b}(X) = \mathrm{CH}^b(X)$, $H^{2b+1,b}(X) = 0$, $H^{2b,b}(X \times X) = \mathrm{CH}^b(X \times X)$ and there is a map

$$H^{2b+2,b}(Z) \rightarrow H^{2b,b}(X \times X \times X) = \mathrm{CH}^b(X \times X \times X)$$

such that the composition

$$\mathrm{CH}^b(X \times X) \xrightarrow{\alpha} H^{2b+2,b}(Z) \rightarrow \mathrm{CH}^b(X \times X \times X)$$

equals $\pi_0^* - \pi_1^* + \pi_2^*$.

Summarizing we see that the invariant μ gives rise to an element $\rho = \rho(\mu)$ in the homology of the complex

$$\mathrm{CH}^b(X) \xrightarrow{\pi_0^* - \pi_1^*} \mathrm{CH}^b(X \times X) \xrightarrow{\pi_0^* - \pi_1^* + \pi_2^*} \mathrm{CH}^b(X \times X \times X).$$

If the element ρ (as a cycle in $\mathrm{CH}^b(X \times X)$) satisfies additionally the condition $c(\rho) \neq 0 \bmod p$, then ρ turns out to be a special correspondence on X .

5 Norm varieties and special correspondences

We continue to assume that $\mathrm{char} k = 0$. The goal of this section is to prove the following theorem:

5.1 Theorem. *Let X be a smooth projective irreducible variety which possesses a special correspondence of type (n, p) and has no zero-cycles of degree coprime to p . Then there exists a (functorial) element $0 \neq u \in H^n(k, \mu_p^{\otimes(n-1)})$ such that X is a norm variety for u .*

For any field extension K/k the invariant $u_K = 0$ iff X_K has a zero cycle of degree coprime to p .

The proof goes as follows:

5.2 Proposition (M. Rost). *Let X be a smooth projective irreducible variety which possesses a special correspondence ρ . Assume $I(X) \subset p\mathbb{Z}$. Then the motive of X considered with $\mathbb{Z}_{(p)}$ -coefficients has an indecomposable generically split direct summand R such that*

$$R \otimes X \simeq \bigoplus_{i=0}^{p-1} X\{bi\}.$$

This Proposition is proved in [Ro07, Proposition 7.14]. The motive R is called a *generalized Rost motive*. Here is an outline of Rost's proof.

The proof of [Ro07, Proposition 7.14] shows that $R = (X, \rho^{p-1}) \bmod p$. By Manin's identity principle (Yoneda lemma for motives) it suffices to construct a correspondence

$$\theta \in \text{Hom}\left(\bigoplus_{i=0}^{p-1} X\{bi\}, X \otimes R\right)$$

such that for any pure motive T

$$\theta \circ -: \text{Hom}\left(T, \bigoplus_{i=0}^{p-1} X\{bi\}\right) \rightarrow \text{Hom}(T, X \otimes R)$$

is an isomorphism. Note that the composition with θ equals the realization of $\Delta_T \otimes \theta$.

Define

$$\theta_i = (\Delta_X \times 1)(1 \times \rho^i) \in \text{Hom}(X\{b(p-1-i)\}, X \otimes R), \quad i = 0, \dots, p-1,$$

and

$$\theta = \sum_{i=0}^{p-1} \theta_i.$$

Let $Y = T \times X$ and $f: Y \rightarrow X$ be the projection. For $0 \leq i \leq p-1$ let

$$\begin{aligned} \varphi_i: \text{CH}^h(Y) &\rightarrow \text{CH}^{h+bi}(Y \times X) \\ \alpha &\mapsto \pi_X^*(\alpha) \cdot (f \times \text{id}_X)^*(\rho^i), \end{aligned}$$

where $\pi_X: Y \times X \rightarrow Y$ is the projection,

$$\Phi = \sum_{i=0}^{p-1} \varphi_i$$

and

$$\begin{aligned} \psi_i: \mathrm{CH}^{h+bi}(Y \times X) &\rightarrow \mathrm{CH}^h(Y) \\ \beta &\mapsto (\pi_X)_*(\beta \cdot (f \times \mathrm{id}_X)^*(\rho^{p-1-i})), \end{aligned}$$

$$\Psi = \sum_{i=0}^{p-1} \psi_i.$$

One verifies using projection formulas that the realization of $\Delta_T \otimes \theta$ equals Φ . In [Ro07, Proof of Proposition 7.14] Rost shows that Φ is an isomorphism. Therefore θ is an isomorphism.

Observe also the following: Let $\overline{X} = X_{k(X)}$ and $\overline{R} = R_{k(X)}$. Then a direct computation shows that over $k(X)$ (where R becomes $\bigoplus_{i=0}^{p-1} \mathbb{Z}/p\{bi\}$) the constructed isomorphism

$$\bigoplus_{i=0}^{p-1} \overline{X}\{bi\} \xrightarrow{\theta_{k(X)}} \overline{X} \otimes \overline{R} = \bigoplus_{i=0}^{p-1} \overline{X}\{bi\}$$

is a lower triangular matrix with $\Delta_{\overline{X}}$ on the diagonal, i.e., $\theta_{k(X)}: \overline{X}\{bi\} \rightarrow \overline{X}\{bj\}$ is zero for $i > j$, and equals $\Delta_{\overline{X}}$ for $i = j$.

From now on we assume that X is an irreducible variety with a special correspondence ρ and with a motive given by Proposition 5.2 and its proof in [Ro07].

5.3 Lemma. *Let X be a smooth projective irreducible variety which possesses a special correspondence ρ . Assume $I(X) \subset p\mathbb{Z}$, and let R be a direct summand of the motive of X (with $\mathbb{Z}_{(p)}$ - or $\mathbb{Z}/p$ -coefficients) given in Proposition 5.2. Then there exists the following filtration of R consisting of exact triangles:*

$$\begin{aligned} R_{p-2}\{b\} &\rightarrow R_{p-1} \rightarrow \mathcal{X} \\ R_{p-3}\{b\} &\rightarrow R_{p-2} \rightarrow \mathcal{X} \\ &\dots \end{aligned}$$

$$R_0\{b\} \rightarrow R_1 \rightarrow \mathcal{X}$$

$$0 \rightarrow R_0 \rightarrow \mathcal{X},$$

where $R_{p-1} = R$ and $\mathcal{X}$ denotes the motive of $\check{C}(X)$.

Proof. We write $\mathbb{Z}/p$ for the coefficient ring. The proof goes by induction.

First of all, note that $\mathcal{X}$ lies in the localizing subcategory generated by X . Let now M be a motive in the localizing subcategory generated by X such that there is an isomorphism $\theta: \bigoplus_{i \in I} X\{i\} \xrightarrow{\sim} M \otimes X$ for some finite set I of non-negative integers. Denote $\overline{X} = X \times_k k(X)$ and assume additionally that there is an isomorphism $\zeta = (\zeta_i)_{i \in I}: M_{k(X)} \rightarrow \bigoplus_{i \in I} \mathbb{Z}/p\{i\}$ defined over $k(X)$ such that the composite morphism

$$\bigoplus_{i \in I} \overline{X}\{i\} \xrightarrow{\theta_{k(X)}} (M \otimes X)_{k(X)} \xrightarrow{\zeta \otimes \text{id}_{\overline{X}}} \bigoplus_{i \in I} \overline{X}\{i\}$$

is a lower triangular matrix with $\Delta_{\overline{X}}$ on the diagonal. In particular, the following diagram commutes:

$$\begin{array}{ccc} (M \otimes X)_{k(X)} & \xrightarrow{\zeta_{\min I} \otimes \text{id}_{\overline{X}}} & \overline{X}\{\min I\} \\ \downarrow \theta_{k(X)}^{-1} & & \downarrow = \\ \bigoplus_{i \in I} \overline{X}\{i\} & \longrightarrow & \overline{X}\{\min I\} \end{array}$$

(the bottom map is the canonical projection).

Observe that these conditions are satisfied by the motive $R = R_{p-1}$ given in the previous proposition. So, we can start induction with $M = R_{p-1}$ and $I = \{bi, i = 0, \dots, p-1\}$. Note also that the isomorphism $\zeta = (\zeta_i)$ is unique up to a sequence of invertible scalars, since $\text{End}(\bigoplus_{i \in I} \mathbb{Z}/p\{i\}) = \bigoplus_{|I|} \mathbb{Z}/p$.

Consider the skeleton filtration of $\mathcal{X}$. Let $Y = \text{Cone}(X \rightarrow \mathcal{X})$ be the cone of the canonical map $X \rightarrow \mathcal{X}$ induced by the structural map $X \rightarrow \text{Spec } k$. Then we have the following exact triangle:

$$M \otimes X \rightarrow M \otimes \mathcal{X} \rightarrow M \otimes Y. \quad (2)$$

By Lemma 3.8(1) $M \otimes \mathcal{X} = M$. Let now l be the smallest element in I . By Lemma 3.1 the above triangle induces the long exact sequence

$$\begin{aligned} \text{Hom}(M \otimes Y, \mathbb{Z}/p\{l\}) &\rightarrow \text{Hom}(M, \mathbb{Z}/p\{l\}) \rightarrow \text{Hom}(M \otimes X, \mathbb{Z}/p\{l\}) \\ &\rightarrow \text{Hom}((M \otimes Y)[-1], \mathbb{Z}/p\{l\}). \end{aligned} \quad (3)$$

The group $\mathrm{Hom}(M \otimes Y, \mathbb{Z}/p\{l\}) = 0$, since $\mathrm{Hom}(V\{l\}[j], \mathbb{Z}/p\{l\}) = \mathrm{Hom}(V[j], \mathbb{Z}/p) = 0$ for any smooth projective variety V and any $j > 0$, and since Y is an extension of $X^i[i-1]$, $i > 1$.

Moreover, we have an injection

$$\mathrm{Hom}((M \otimes Y)[-1], \mathbb{Z}/p\{l\}) \rightarrow \mathrm{Hom}(M \otimes X \otimes X, \mathbb{Z}/p\{l\})$$

induced by the natural map $(X \otimes X)[1] \rightarrow Y$. Observe that the composition $(X \otimes X)[1] \rightarrow Y \rightarrow X[1]$ is the difference of two projections $\pi_0 - \pi_1$ (cf. Section 4 and [Vi98, p. 31]).

Since by induction hypothesis $M \otimes X \simeq \bigoplus_{i \in I} X\{i\}$ and since l is minimal, we have $\mathrm{Hom}(M \otimes X, \mathbb{Z}/p\{l\}) \simeq \mathrm{Hom}(X\{l\}, \mathbb{Z}/p\{l\}) = \mathrm{Hom}(X, \mathbb{Z}/p) = \mathrm{Ch}^0(X)$.

We claim next that the image of the structural map $\mathrm{pr}: X \rightarrow \mathrm{Spec} k$ (as an element of $\mathrm{Hom}(M \otimes X, \mathbb{Z}/p\{l\})$) in $\mathrm{Hom}((M \otimes Y)[-1], \mathbb{Z}/p\{l\})$ is trivial. Indeed, since $\mathrm{Hom}(X, \mathbb{Z}/p)$ injects into $\mathrm{Hom}(X_{k(X)}, \mathbb{Z}/p)$, it suffices to show that the image of $\mathrm{pr}_{k(X)}$ in $\mathrm{Hom}((M \otimes X \otimes X)_{k(X)}, \mathbb{Z}/p\{l\})$ is zero. But this is clear, since over $k(X)$ the motive M is a direct sum of Tate motives. Therefore, since sequence (3) is exact, we can find a preimage $\beta \in \mathrm{Hom}(M, \mathbb{Z}/p\{l\})$ of the structural map.

Lemma 3.8(2) gives us a map $M \rightarrow \mathcal{X}\{l\}$. Define

$$M' = \mathrm{Cone}(M \rightarrow \mathcal{X}\{l\})[-1]$$

and consider the map

$$\gamma: M' \otimes X \rightarrow M \otimes X \rightarrow \bigoplus_{i \in I \setminus \{l\}} X\{i\}$$

arising from the exact triangle $(M' \rightarrow M \rightarrow \mathcal{X}\{l\}) \otimes X$ and the isomorphism $\theta^{-1}: M \otimes X \simeq \bigoplus_{i \in I} X\{i\}$. By our inductive assumptions on M one can see that $\gamma_{k(X)}$ is an isomorphism. Therefore by [Vi98, Theorem 2.3.5] we have an isomorphism $M' \otimes X \otimes X \simeq \bigoplus_{i \in I \setminus \{l\}} X \otimes X\{i\}$. So, we can replace X by $X \otimes X$ and apply induction. Note that the motives of $\check{C}(X)$ and of $\check{C}(X \times X)$ are isomorphic by [Vi98, Theorem 2.3.4], and γ satisfies all hypothesis needed for the next inductive step.

Therefore by induction we can construct the motives

$$M_{p-1} := R, M_{p-2}, M_{p-3}, \dots, M_0, M_{-1}$$

together with exact triangles

$$M_{s-1} \rightarrow M_s \rightarrow \mathcal{X}\{b(p-1-s)\}, \quad s = 0, \dots, p-1,$$

and such that $M_s \otimes X^{p-s} \simeq \bigoplus_{i=p-1-s}^{p-1} X^{p-s}\{bi\}$ for all s . In particular, $M_{-1} \otimes X^{p+1} = 0$.

But then $M_{-1} = M_{-1} \otimes \mathcal{X} = 0$. Finally, the existence of exact triangles as in the statement of the lemma follows from the cancellation theorem (see Lemma 3.6). We are done. $\square$

Next we investigate the non-triviality of the construction of Lemma 5.3.

5.4 Lemma. *For $i = 1, \dots, p-1$ if $c_1 > c_2$ then*

$$\mathrm{Hom}(R_i(c_1)[d_1], \mathbb{Z}/p(c_2)[d_2]) = 0.$$

Proof. The statement follows from Lemma 5.3 and Lemma 3.12. $\square$

5.5 Lemma. *Let X be a variety over k which possesses a special correspondence ρ of type (n, p) . Assume that $I(X) \subset p\mathbb{Z}$. Let K/k be a field extension. Then $\mathrm{Hom}(\mathcal{X}_K, \mathbb{Z}/p\{b\}[1]) = 0$ iff X_K has a zero-cycle of degree coprime to p .*

Proof. Consider Hom from the exact triangle $R_{p-2}\{b\} \rightarrow R_{p-1} \rightarrow \mathcal{X}$ to $\mathbb{Z}/p\{b\}$. We get the following long exact sequence:

$$\begin{aligned} \mathrm{Hom}(R_{p-1}, \mathbb{Z}/p\{b\}) &\rightarrow \mathrm{Hom}(R_{p-2}\{b\}, \mathbb{Z}/p\{b\}) \\ &\rightarrow \mathrm{Hom}(\mathcal{X}[-1], \mathbb{Z}/p\{b\}) \rightarrow \mathrm{Hom}(R_{p-1}[-1], \mathbb{Z}/p\{b\}). \end{aligned}$$

Note that $\mathrm{Hom}(R_{p-1}, \mathbb{Z}/p\{b\}) = \mathrm{Ch}^b(R_{p-1})$ (Lemma 3.3), since R_{p-1} is a pure motive, $\mathrm{Hom}(R_{p-2}\{b\}, \mathbb{Z}/p\{b\}) = \mathrm{Hom}(R_{p-2}, \mathbb{Z}/p)$ (Lemma 3.6), $\mathrm{Hom}(\mathcal{X}[-1], \mathbb{Z}/p\{b\}) = \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p\{b\}[1])$, and $\mathrm{Hom}(R_{p-1}[-1], \mathbb{Z}/p\{b\}) = 0$ (Lemma 3.3).

Moreover, it follows from the proof of Lemma 5.3 that the map

$$\mathrm{Hom}(R_{p-2}, \mathbb{Z}/p) \rightarrow \mathrm{Hom}(X, \mathbb{Z}/p) = \mathrm{Ch}^0(X)$$

coming from triangle (2) is an isomorphism. Thus, we get the following commutative diagram

$$\begin{array}{ccccccc}
\mathrm{Hom}(R_{p-1}, \mathbb{Z}/p\{b\}) & \longrightarrow & \mathrm{Hom}(R_{p-2}\{b\}, \mathbb{Z}/p\{b\}) & \xrightarrow{g} & \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p\{b\}[1]) & \longrightarrow & 0 \\
\parallel & & \downarrow \simeq & & \parallel & & \\
\mathrm{Ch}^b(R_{p-1}) & \xrightarrow{f} & \mathrm{Ch}^b(X\{b\}) & \xrightarrow{g} & \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p\{b\}[1]) & \longrightarrow & 0
\end{array}$$

Moreover, we have an injection $\mathrm{Ch}^b(X\{b\}) \rightarrow \mathrm{Ch}^b(X_{k(X)}\{b\})$. On the other hand, the image of $\mathrm{CH}^b(R_{p-1}) \rightarrow \mathrm{CH}^b((R_{p-1})_{k(X)}) = \mathbb{Z}$ is $p\mathbb{Z}$ if X has no zero-cycles of degree coprime to p or $\mathbb{Z}$ otherwise (see [Ro07, Corollary 5.7]). This implies the lemma. $\square$

5.6 Lemma. *Define $\mu \in \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p\{b\}[1])$ as the image of $1 = [X] \in \mathrm{Ch}^0(X)$ under the map g given in the proof of Lemma 5.5. Then for $0 \leq i \leq n-2$ we have $Q_i(\mu) = 0$.*

Proof. Since μ is defined as the image of some element in $\mathrm{Hom}(R_{p-2}, \mathbb{Z}/p)$ and the Milnor operations respect pull-backs, it suffices to show that

$$Q_i(\mathrm{Hom}(R_{p-2}, \mathbb{Z}/p)) = 0.$$

The exact triangle $R_{p-1} \rightarrow \mathcal{X} \rightarrow R_{p-2}\{b\}[1]$ gives the exact sequence

$$\begin{aligned}
\mathrm{Hom}(R_{p-2}(b)[2b+1], \mathbb{Z}/p(p^i-1)[2p^i-1]) &\rightarrow \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p(p^i-1)[2p^i-1]) \\
&\rightarrow \mathrm{Hom}(R_{p-1}, \mathbb{Z}/p(p^i-1)[2p^i-1]).
\end{aligned}$$

The first group in this sequence is zero by Lemma 5.4, since $b > p^i - 1$ by our assumptions. The last group is zero, since R_{p-1} is a pure motive and $2p^i - 1 > 2(p^i - 1)$. Therefore $\mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p(p^i-1)[2p^i-1]) = 0$.

The exact triangle $R_{p-3}\{b\} \rightarrow R_{p-2} \rightarrow \mathcal{X}$ gives the exact sequence

$$\begin{aligned}
\mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p(p^i-1)[2p^i-1]) &\rightarrow \mathrm{Hom}(R_{p-2}, \mathbb{Z}/p(p^i-1)[2p^i-1]) \\
&\rightarrow \mathrm{Hom}(R_{p-3}\{b\}, \mathbb{Z}/p(p^i-1)[2p^i-1]).
\end{aligned}$$

The first and the last groups are zero by Lemma 5.4 and by the previous considerations. The lemma is proved. $\square$

The following lemma is due to V. Voevodsky. We reproduce its proof for reader's convenience.

5.7 Lemma ([Vo03, Proof of Lemma 6.7]). *Let $\delta \in H^{n,n-1}(\mathcal{X}, \mathbb{Z}/p)$. If*

$$Q_{n-2} \dots Q_1 Q_0(\delta) = 0$$

then $\delta = 0$.

Proof. The exact triangle of pointed simplicial schemes

$$\mathcal{X}_+ \rightarrow \operatorname{Spec} k_+ \rightarrow \tilde{\mathcal{X}}$$

defines a monomorphism $H^{d,c}(\mathcal{X}, \mathbb{Z}/p) \rightarrow \tilde{H}^{d+1,c}(\tilde{\mathcal{X}}, \mathbb{Z}/p)$ for any $d > c$. Thus, we can identify δ with its image $\tilde{\delta}$ in $\tilde{H}^{n+1,n-1}(\tilde{\mathcal{X}}, \mathbb{Z}/p)$.

Assume that $\tilde{\delta} \neq 0$. We want to show that $Q_i Q_{i-1} \dots Q_0(\tilde{\delta}) \neq 0$ for all $i \leq n-2$. Assume by induction that $Q_{i-1} \dots Q_0(\tilde{\delta}) \neq 0$. If $Q_i Q_{i-1} \dots Q_0(\tilde{\delta}) = 0$, then by Lemma 3.11 there exists v such that $Q_i(v) = Q_{i-1} \dots Q_0(\tilde{\delta})$.

A straightforward computation shows that $v \in \tilde{H}^{d,c}(\tilde{\mathcal{X}}, \mathbb{Z}/p)$ where $c = n - i - p^i + 1 + \frac{p^i - p}{p-1}$ and $d = n - i + 2 - 2p^i + 2 \cdot \frac{p^i - p}{p-1}$.

But $d - c \leq 1$. Therefore by Lemma 3.14 $v = 0$. Contradiction. $\square$

5.8 Lemma. *Assume that $p > 2$. Then in the above notation there exists $\delta \in \operatorname{Hom}(\mathcal{X}, \mathbb{Z}/p(n-1)[n])$ such that $\mu = Q_{n-2} \dots Q_1 Q_0(\delta)$.*

Proof. As in the previous lemma it suffices to prove this identity in the reduced motivic cohomology of $\tilde{\mathcal{X}}$. Let $\eta \in \tilde{H}^{2b+2,b}(\tilde{\mathcal{X}}, \mathbb{Z}/p)$ be the image of μ .

Assume we have constructed a sequence of elements $\eta_{n-2}, \eta_{n-3}, \dots, \eta_{n-s}$ for some $2 \leq s < n$ such that $\eta_{n-i+1} = Q_{n-i}(\eta_{n-i})$ and

$$\eta = Q_{n-2} Q_{n-3} \dots Q_{n-s}(\eta_{n-s}).$$

For $s = 2$ such an element η_{n-2} with $\eta = Q_{n-2}(\eta_{n-2})$ exists by Lemma 3.11, since $Q_{n-2}(\eta) = 0$ by Lemma 5.6.

We claim that there exists η_{n-s-1} satisfying the same conditions. To prove this it suffices to show that $v := Q_{n-s-1}(\eta_{n-s})$ equals 0.

Consider

$$\begin{aligned} Q_{n-2} Q_{n-3} \dots Q_{n-s}(v) &= Q_{n-2} Q_{n-3} \dots Q_{n-s} Q_{n-s-1}(\eta_{n-s}) \\ &= \pm Q_{n-s-1}(Q_{n-2} Q_{n-3} \dots Q_{n-s}(\eta_{n-s})) = \pm Q_{n-s-1}(\eta) = 0, \end{aligned}$$

since $0 \leq n - s - 1 \leq n - 3$.

A straightforward computation shows that

$$\eta_{n-s} \in \tilde{H}^{2b+s-2, \frac{p^{n-1}-p^{n-s}}{p-1}+1, b+s-1-\frac{p^{n-1}-p^{n-s}}{p-1}}(\tilde{\mathcal{X}}, \mathbb{Z}/p)$$

and

$$v \in \tilde{H}^{2b+s-2, \frac{p^{n-1}-p^{n-s}}{p-1}+2p^{n-s-1}, b+s-2-\frac{p^{n-1}-p^{n-s}}{p-1}+p^{n-s-1}}(\tilde{\mathcal{X}}, \mathbb{Z}/p).$$

Let $2 \leq t \leq s$. We claim that if $Q_{n-t}(Q_{n-t-1} \dots Q_{n-s}(v)) = 0$, then $Q_{n-t-1} \dots Q_{n-s}(v) = 0$. Indeed, if $Q_{n-t}(Q_{n-t-1} \dots Q_{n-s}(v)) = 0$, then there exists w such that $Q_{n-t}(w) = Q_{n-t-1} \dots Q_{n-s}(v)$. A straightforward computation shows that $w \in \tilde{H}^{d,c}(\tilde{\mathcal{X}}, \mathbb{Z}/p)$, where

$$c = \frac{p^{n-t}-1}{p-1} - 1 + p^{n-s-1} + t - p^{n-t}$$

and

$$d = 2 \cdot \frac{p^{n-t}-1}{p-1} + 2p^{n-s-1} + t - 2p^{n-t} + 1.$$

Another straightforward computation shows that $d \leq c + 1$ if $p > 2$, and by Lemma 3.14 $w = 0$. Therefore $v = 0$. $\square$

Proof of Theorem 5.1 for $p > 2$: The element δ constructed in Lemma 5.8 can be identified with an element $u \in H_{et}^n(k, \mu_p^{\otimes(n-1)})$ by Lemma 3.15. It remains to show that for any field extension K/k

$$u_K = 0 \text{ iff } X_K \text{ has a zero-cycle of degree coprime to } p.$$

By construction $u_K = 0$ iff $\delta_K = 0$. By Lemma 5.7 $\delta_K = 0$ iff $\mu_K = 0$. Finally by Lemma 5.5 $\mu_K = 0$ iff X_K has a zero-cycle of degree coprime to p . The theorem is proved.

6 Binary motives

In this section we determine the structure of binary direct summands of smooth projective varieties.

6.1 Theorem. *Let X be a smooth projective irreducible variety over a field k with no zero-cycles of odd degree. Let M be a (binary) direct summand of X with $M \otimes X \simeq X \oplus X\{\dim X\}$. Then*

1. $\dim X = 2^{n-1} - 1$ for some n .
2. *There exists a (functorial) element $u \in H^n(k, \mathbb{Z}/2)$ such that for any field extension K/k we have $u_K = 0$ iff X_K has a zero-cycle of odd degree.*

One can notice that Rost's proof ([Ro07, Proof of Theorem 9.1 and Lemma 9.10]) implies the following lemma. For reader's convenience we sketch Rost's proof below.

6.2 Lemma. *Let X be a smooth projective irreducible variety of dimension d with no zero-cycles of odd degree which possesses a cycle $r \in \text{CH}_d(X \times X)$ such that $r_{k(X)} \bmod 2$ is a projector and $(X_{k(X)}, r_{k(X)}) \simeq \mathbb{Z}/2 \oplus \mathbb{Z}/2\{d\}$.*

Then $d = 2^{n-1} - 1$ for some n and X is a ν_{n-1} -variety.

Proof. Let $S_\bullet$ (resp. $S^\bullet$) denote the homological (resp. cohomological) Steenrod operations in the Chow theory (see [Br03a]).

Then $S_\bullet(\alpha) = S^\bullet(\alpha)b_\bullet(X)$, $\alpha \in \text{Ch}(X)$, where $b_\bullet$ is the (total) characteristic number defined in 2.2 for partition $(p-1, \dots, p-1)$ (Since $p = 2$, we have $b_\bullet(X) = c_\bullet(-T_X)$).

One has $b_d(X) = S_d([X])$. By our assumptions the cycle r has the property $(\pi_0)_*(r) = c(r)[X]$ with $c(r) = 1 \bmod 2$. Moreover, $S_\bullet(r) = S^\bullet(r)b_\bullet(X^2)$.

We show first that $b_d(X) \neq 0 \bmod p^2$. To do this it suffices to find an integral representation of the cycle $S_d(r)$ with degree not divisible by 4.

We have

$$S_d(r) = \sum_{i=0}^d S^i(r)b_{d-i}(X^2). \quad (4)$$

The last term $S^d(r) = r^2 \bmod 2$ and $\deg r^2 = 2c(r)^2$ is not divisible by 4. Therefore it suffices to show that all other terms of this sum have integral representatives whose degrees are divisible by 4.

Let F be the function field of X . For the first term of (4) we have

$$r_F b_d(X^2) = H \times b_d(X) + b_d(X) \times H,$$

where H is the class of a rational point on X_F . This cycle has degree divisible by 4.

Let π be an integral representative of $S^i(r)$. We want to show that $\deg(\pi b_{d-i}(X^2))$ is divisible by 4. By dimension reasons $S^i(H) = 0$ for $1 \leq i \leq d-1$. Therefore $S^i(r_F) = 0$ for such i 's. Thus, there exists a cycle γ over F such that $\pi_F = 2\gamma$. It remains to show that $\deg(\gamma b_{d-i}(X^2))$ is divisible by 2.

We set $X = X_0 = X_1$ and write $X^2 = X_0 \times X_1$. We have

$$b_{d-i}(X^2) = \sum_{j+r=d-i} b_j(X_0)b_r(X_1).$$

By assumptions $j+r = d-i > 0$. Therefore $r > 0$ or $j > 0$. In the first case $\deg(\gamma b_j(X_0)b_r(X_1)) = \deg((\pi_0)_*(\gamma b_j(X_0)b_r(X_1)))$. Therefore it suffices to show that $\deg(\alpha_F \beta)$ is divisible by 2 for all $\alpha \in \text{CH}^r(X)$, $\beta \in \text{CH}^{d-r}(X_F)$.

Let φ be a preimage of β under the natural map

$$\mathrm{CH}^{d-r}(X^2) \rightarrow \mathrm{CH}^{d-r}(X \times \mathrm{Spec} F).$$

Consider the cycle $\omega = r(\alpha \times 1)\varphi \in \mathrm{CH}_0(X^2)$.

One has $\omega_F = (H \times 1)(\alpha_F \times 1)\varphi_F + (1 \times H)(\alpha_F \times 1)\varphi_F = (1 \times H)(\alpha_F \times 1)\varphi_F$. Therefore $\deg \omega = \deg(\alpha_F \beta) = (\pi_0)_*((\alpha_F \times 1)\varphi_F) = c(r) \deg(\alpha_F \beta)$ is divisible by 2, since X^2 has no zero-cycles of odd degree.

The case $j > 0$ is similar.

Thus $b_d(X) \neq 0 \bmod p^2$. By [Ro07, Lemma 9.10] $\dim X = 2^{n-1} - 1$ for some n . Finally, by [Ro07, Lemma 9.13] X is a ν_{n-1} -variety. $\square$

By Lemma 5.3 we have an exact triangle

$$\mathcal{X}\{b\} \rightarrow M \rightarrow \mathcal{X} \xrightarrow{\mu'} \mathcal{X}\{b\}[1].$$

Denote $\mu \in \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/2(b)[2b+1])$ the pull-back of μ' under the map of Lemma 3.8(2). Observe that the construction of μ here is compatible with the construction of μ of Section 5 for $p = 2$.

The following lemmas were proven by O. Izhboldin and A. Vishik in the case when X is a quadric. Nevertheless their proofs are general and do not use any specific of quadrics. For reader's convenience we reproduce them below.

6.3 Lemma ([IzhV00, Sublemma 6.3]). *The map*

$$\mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p(c)[d]) \rightarrow \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/p(c+b)[d+2b+1])$$

induced by the multiplication by μ is surjective for $d \geq c$.

Proof. Consider the morphism from the exact triangle

$$M \rightarrow \mathcal{X} \xrightarrow{\mu'} \mathcal{X}(b)[2b+1]$$

to $\mathbb{Z}/2(b+c)[2b+d+1]$.

We get the long exact sequence

$$\begin{aligned} \mathrm{Hom}(\mathcal{X}(b)[2b+1], \mathbb{Z}/2(b+c)[2b+d+1]) &\xrightarrow{\mu'^*} \mathrm{Hom}(\mathcal{X}, \mathbb{Z}/2(b+c)[2b+d+1]) \\ &\rightarrow \mathrm{Hom}(M, \mathbb{Z}/2(b+c)[2b+d+1]). \end{aligned}$$

Note that $\text{Hom}(\mathcal{X}(b)[2b+1], \mathbb{Z}/2(b+c)[2b+d+1]) = \text{Hom}(\mathcal{X}, \mathbb{Z}/2(c)[d])$ and by Lemma 3.13 the map μ'^* coincides with the multiplication by μ . Moreover, $\text{Hom}(M, \mathbb{Z}/2(b+c)[2b+d+1]) = 0$ by Lemma 3.5, since M is a pure motive and $2b+d+1 - (b+c) = b+d-c+1 > b = \dim X$.

Therefore the multiplication by μ is surjective for $d \geq c$. $\square$

Let $\tilde{\mathcal{X}} = \text{Cone}(\mathcal{X} \rightarrow \text{Spec } k)$.

6.4 Lemma ([IzhV00, Sublemma 6.6]). *For any $i = 0, 1, \dots, n-2$ the Milnor operation*

$$Q_i: \tilde{H}^{d,c}(\tilde{\mathcal{X}}, \mathbb{Z}/2) \rightarrow \tilde{H}^{d+2^{i+1}-1, c+2^i-1}(\tilde{\mathcal{X}}, \mathbb{Z}/2)$$

is injective if $d - c = b + 2 + 2^i$.

Proof. Let $v \in \tilde{H}^{d,c}(\tilde{\mathcal{X}}, \mathbb{Z}/2)$ with $Q_i(v) = 0$. We want to show that then $v = 0$.

Since $Q_i(v) = 0$, by Lemma 3.11 there exists

$$t \in \tilde{H}^{d-2^{i+1}+1, c-2^i+1}(\tilde{\mathcal{X}}, \mathbb{Z}/2)$$

such that $v = Q_i(t)$.

On the other hand, the exact triangle of pointed simplicial schemes

$$\tilde{\mathcal{X}}[-1] \xrightarrow{\tau} \mathcal{X}_+ \rightarrow \text{Spec } k_+$$

induces an isomorphism

$$\tau^*: H^{d-2^{i+1}+1, c-2^i+1}(\mathcal{X}, \mathbb{Z}/2) \rightarrow \tilde{H}^{d-2^{i+1}+2, c-2^i+1}(\tilde{\mathcal{X}}, \mathbb{Z}/2).$$

Therefore there exists

$$w \in \text{Hom}(\mathcal{X}, \mathbb{Z}/2(c - 2^i + 1)[d - 2^{i+1} + 1])$$

with $\tau^*(w) = t$. Moreover, by Lemma 6.3 there exists

$$u \in \text{Hom}(\mathcal{X}, \mathbb{Z}/2(c - 2^i + 1 - b)[d - 2^{i+1} - 2b])$$

such that $w = \mu \cdot u$.

By Lemma 3.9(3) we have $Q_i(\mu \cdot u) = Q_i(\mu) \cdot u + \mu \cdot Q_i(u) + \sum \xi^{n_j} \varphi_j(\mu) \psi_j(u)$, where ξ is the class of -1 in $\text{Hom}(\mathbb{Z}/2, \mathbb{Z}/2(1)[1])$, $n_j > 0$ and φ_j, ψ_j are some homogeneous cohomological operations of some bidegrees $(*)[*']$ with $*' > 2* \geq 0$.

Since $e := d - 2^{i+1} - 2b = c - 2^i + 1 - b$ by assumptions, Lemma 3.14 implies an isomorphism $\text{pr}^*: \text{Hom}(\mathbb{Z}/2, \mathbb{Z}/2(e)[e]) \rightarrow \text{Hom}(\mathcal{X}, \mathbb{Z}/2(e)[e])$, where $\text{pr}: \mathcal{X} \rightarrow \text{Spec } k$ is the structural map. Therefore there exists

$$u_0 \in \text{Hom}(\mathbb{Z}/2, \mathbb{Z}/2(e)[e])$$

with $u = \text{pr}^*(u_0)$.

On the other hand, since $\text{Hom}(\mathbb{Z}/2, \mathbb{Z}/2(f)[g]) = 0$ for any $g > f$, we have $Q_i(u_0) = 0$ and $\psi_j(u_0) = 0$. Since the Milnor operations commute with pull-backs, we also have $Q_i(u) = 0$ and $\psi_j(u) = 0$.

Summarizing we obtain:

$$v = Q_i(t) = Q_i(\tau^*(w)) = \tau^*(Q_i(w)) = \tau^*(Q_i(\mu \cdot u)) = \tau^*(Q_i(\mu) \cdot u) = 0$$

the last equality by Lemma 5.6. $\square$

6.5 Lemma ([IzhV00, Sublemma 6.7]). *Lemma 5.8 holds for $p = 2$, i.e., there exists $\delta \in \text{Hom}(\mathcal{X}, \mathbb{Z}/2(n-1)[n])$ such that $\mu = Q_{n-2} \dots Q_1 Q_0(\delta)$.*

Proof. As in the proofs of Lemmas 5.7 and 5.8 we identify μ with its image in the reduced motivic cohomology $\tilde{H}^{2b+2,b}(\mathcal{X}, \mathbb{Z}/p)$ which we denote as $\eta = \eta_{-1}$.

Assume by induction that we have constructed an element η_m , $-1 \leq m \leq n-3$, such that $\eta = Q_m \dots Q_1 Q_0(\eta_m)$. We want to show that there exists η_{m+1} with $\eta_m = Q_{m+1}(\eta_{m+1})$.

By Lemma 3.11 it suffices to show that $v := Q_{m+1}(\eta_m) = 0$. We have $Q_m Q_{m-1} \dots Q_0(v) = Q_m Q_{m-1} \dots Q_0 Q_{m+1}(\eta_m) = Q_{m+1} Q_m Q_{m-1} \dots Q_0(\eta_m) = Q_{m+1}(\eta) = 0$ by Lemma 5.6.

A straightforward computation shows that for any $0 \leq t \leq m$ the element $Q_{t-1} \dots Q_0(v) \in \tilde{H}^{d,c}(\mathcal{X}, \mathbb{Z}/2)$ with $d - c = b + 2 + 2^t$. By Lemma 6.4 Q_t is injective on $\tilde{H}^{d,c}(\mathcal{X}, \mathbb{Z}/2)$ for such c and d . Therefore the equality $Q_m Q_{m-1} \dots Q_0(v) = 0$ implies $v = 0$. $\square$

Proof of Theorem 6.1: The first part of the Theorem follows from Lemma 6.2.

The proof of the second assertions repeats word by word the proof of Theorem 5.1 given at the end of the previous section with Lemma 5.8 replaced by Lemma 6.5.

Proof of Theorem 5.1 for $p = 2$: This is a particular case of Theorem 6.1.

7 Compression of varieties

The goal of this section is to provide a certain compression algorithm for smooth projective varieties. We assume that $\text{char } k = 0$.

7.1 Lemma. *Let X be a smooth projective irreducible variety and Y a closed subvariety of X of minimal dimension such that $Y_{k(X)}$ has a zero-cycle of degree coprime to p .*

Then there exists a smooth projective variety $\tilde{Y}$ which is birational to Y such that $\tilde{Y}_{k(X)}$ has a zero-cycle of degree coprime to p .

Proof. Let $U \subset Y$ be the open subvariety of smooth points in Y and $V = Y \setminus U$ the singular locus. We claim that there is a zero-cycle of degree coprime to p supported in $U_{k(X)}$. Indeed, assume the contrary. Then all zero-cycles supported on $U_{k(X)}$ have degrees divisible by p . Since there exists a zero-cycle on $Y_{k(X)}$ of degree coprime to p , the variety $V_{k(X)} \subset X_{k(X)}$ has a zero-cycle of degree coprime to p . Since $\dim V < \dim Y$ and $\dim Y$ is minimal by our assumptions, we come to a contradiction.

Using the standard algorithm of resolution of singularities due to Hironaka we can find a variety $\tilde{Y}$ birational to Y which contains U . This $\tilde{Y}$ has a zero-cycle of degree 1 mod p over $k(X)$. $\square$

7.2 Corollary. *Let X be a generically split smooth projective irreducible k -variety. Let Y be a closed subvariety of X of minimal dimension such that the class $[Y_{k(X)}]$ in $\text{Ch}(X_{k(X)})$ is non-zero. Then there exists a smooth projective variety $\tilde{Y}$ birational to Y such that $\tilde{Y}_{k(X)}$ has a zero-cycle of degree 1 mod p .*

Proof. Since X is generically split, by [KM06, Remark 5.6] there exists a closed subvariety $Z \subset X_{k(X)}$ such that $[Y_{k(X)}] \cdot [Z]$ has degree coprime to p . Since the product $Y_{k(X)} \cdot Z$ in the Chow ring can be represented by a cycle on the intersection $Y_{k(X)} \cap Z$, the variety $Y_{k(X)} \cap Z \subset Y_{k(X)}$ has a zero-cycle of degree coprime to p .

By Lemma 7.1 it suffices to show that Y has minimal dimension among all closed subvarieties of X such that $Y_{k(X)}$ has a zero-cycle of degree coprime to p .

The proof of this claim copies the proof of [KM06, Theorem 5.8].

Let $Y' \xrightarrow{\text{in}} X$ be a closed subvariety of X such that $Y'_{k(X)}$ has a zero-cycle of degree coprime to p . It remains to show that $\dim Y' \geq \dim Y$.

Let $\overline{\text{Ch}}$ denote the image of the restriction map $\text{Ch}(-) \rightarrow \text{Ch}(-_{k(-)})$. It suffices to show that $\overline{\text{Ch}}_i(X)$ is non-zero for some $i \leq \dim Y'$. By [KM06, Corollary 5.4] it suffices to show that the push-forward

$$(in \times \text{id}_X)_*: \overline{\text{Ch}}(Y' \times X) \rightarrow \overline{\text{Ch}}(X \times X)$$

is non-zero.

Denote $F = k(X)$. By assumptions there exists a p -coprime field extension K/F such that $k(Y') \hookrightarrow K$. Let W be the closure of the image of the morphism $K \rightarrow Y' \times X$. Then the cycle $(in \times \text{id}_X)_*([W])$ is non-zero, since for the second projection $\pi_1: X^2 \rightarrow X$ we have

$$(\pi_1)_*(in \times \text{id}_X)_*([W]) = [K : k(X)] \cdot 1 \neq 0.$$

□

7.3 Definition. Let X be a smooth projective variety and $R = (X, \pi)$ a direct summand of the motive of X . We say that R supports 0-cycles of X if $\deg(\pi_*(\text{CH}_0(X))) \neq 0$ (π_* denotes the realization of π).

Finally, we need the following lemma.

7.4 Lemma. *Let X and Y be smooth projective irreducible varieties. Consider the category of Chow motives with $\mathbb{Z}/p$ -coefficients. Assume that X has a zero-cycle of degree 1 mod p over $k(Y)$ and Y has a zero-cycle of degree 1 mod p over $k(X)$. Let R be an indecomposable generically split direct summand of the motive of X which supports zero-cycles. Then R is an indecomposable direct summand of Y which supports zero-cycles.*

Proof. Since X has a zero-cycle of degree 1 mod p over $k(Y)$, there exists by [PSZ, Lemma 1.6] a cycle $\varphi_1 \in \text{Ch}_{\dim X}(X \times Y)$ such that $\varphi_1: pt_X \mapsto pt_Y$ over $k(Y)$.

By symmetry there exists a cycle $\psi_1 \in \text{Ch}_{\dim Y}(Y \times X)$ such that $\psi_1: pt_Y \mapsto pt_X$ over $k(X)$.

Let r be a projector defining R , i.e., $R = (X, r)$. Consider the cycles $\varphi_2 = \varphi_1 \circ r$ and $\psi_2 = r \circ \psi_1$. The composition $\psi_2 \circ \varphi_2 \in \text{End}(R)$ and is non-zero, since $\psi_2 \circ \varphi_2: pt_X \mapsto pt_X$ over $k(X)$.

Consider $\text{End}(R_{k(X)})$. Since R is generically split and indecomposable, it follows that the image of the natural map $\text{End}(R) \rightarrow \text{End}(R_{k(X)})$ equals $\mathbb{Z}/p \cdot r_{k(X)}$. Therefore the cycle $(\psi_2 \circ \varphi_2)_{k(X)}$ equals $c \cdot r_{k(X)}$ with $c \in (\mathbb{Z}/p)^\times$. Multiplying ψ_2 or φ_2 by c^{-1} we may assume that $c = 1$.

By Lemma 3.18 Rost nilpotence holds for R . Therefore $\psi_2 \circ \varphi_2 = r + n$, where n is some nilpotent element in $\text{End}(R)$. Since n is nilpotent, $r + n$ is invertible and $(r + n)^{-1} \circ \psi_2 \circ \varphi_2 = r$. Define now $\psi_3 = (r + n)^{-1} \circ \psi_2$. Then $\psi_3 \circ \varphi_2 = r$, and therefore $\varphi_2 \circ \psi_3$ is a projector on Y and $(Y, \varphi_2 \circ \psi_3) \simeq (X, r) = R$ with mutually inverse isomorphisms φ_2 and ψ_3 . $\square$

8 Serre's question about E_8

Let G be a simple adjoint linear algebraic group of inner type over k with $\text{char } k = 0$. We briefly sketch the definitions of two invariants of G : the J -invariant and the Rost invariant. See [PSZ] and [Inv, §31] for formal definitions.

8.1 (J -invariant). Let p be a prime number.

8.2 Definition. Let R be a motive which is a finite direct sum of Tate motives. The polynomial

$$P(R, t) = \sum_{i \geq 0} a_i t^i \in \mathbb{Z}[t],$$

where $a_i = \dim \text{Ch}_i(R)$ is called the Poincaré polynomial of $R \bmod p$.

Informally speaking, the J -invariant of a simple group G is a discrete invariant which measures the motivic decomposition of the variety of Borel subgroups of G .

Consider the motive of the variety of complete G -flags (variety of Borel subgroups) with $\mathbb{Z}/p$ - or $\mathbb{Z}_{(p)}$ -coefficients. It turns out that it is isomorphic to a direct sum of shifted copies of the same indecomposable generically split motive $R = R_p(G)$ such that over any splitting field K of G the Poincaré polynomial of R_K equals

$$P(R_K, t) = \prod_{i=1}^r \frac{t^{d_i p^{j_i}} - 1}{t^{d_i} - 1} \quad (5)$$

for some given numbers r and $d_1 \leq d_2 \leq \dots \leq d_r$ which depend only on the Dynkin type of G and some r -tuple $(j_1, \dots, j_r)$ of non-negative integers (see [PSZ, Definition 4.5 and Theorem 4.9]).

The tuple $J_p(G) = (j_1, \dots, j_r)$ arising from this decomposition is called the *J-invariant* of G modulo p . We leave the problem of correctness of such a definition aside and refer to [PSZ] for more details.

An important application of the *J-invariant* which is essentially used in the proof of Serre's problem described below is the classification of generically split projective homogeneous varieties given in [PS10, Theorems 5.5 and 5.7].

Note also that formula (5) implies that the variety of Borel subgroups of G has a binary direct summand if $J_2(G) = (0, \dots, 0, 1, 0, \dots, 0)$. In Lemma 8.6 we'll provide a concrete example.

8.3 Example. If G has type E_8 and $p = 2$, then $r = 4$, $d_1 = 3$, $d_2 = 5$, $d_3 = 9$, and $d_4 = 15$. Moreover, $j_4 = 0$ or 1 , and if $j_1 = 0$ then $j_2 = j_3 = 0$ (see [PSZ, §6, Table]).

8.4 (Rost invariant). Let G be a simply connected simple algebraic group over k . The Rost invariant of G is an invariant of G -torsors which takes values in $H^3(k, \mathbb{Q}/\mathbb{Z}(2))$.

Formally: consider the abelian group $\text{Inv}^3(G, \mathbb{Q}/\mathbb{Z}(2))$ of natural transformations of functors $H^1(-, G) \rightarrow H^3(-, \mathbb{Q}/\mathbb{Z}(2))$ defined on the category of field extensions of k .

It turns out that it is a finite cyclic group with a canonical generator called the *Rost invariant* of G . If G is adjoint and simply-connected, then one can identify $H^1(k, G)$ with the (pointed) set of isomorphism classes of the twisted forms of G over k . In particular, in this case one can associate with each twisted form of G an element in $H^3(k, \mathbb{Q}/\mathbb{Z}(2))$ (the Rost invariant).

8.5 Example. Let G be a group of type E_8 . It is well-known that it is adjoint and simply-connected, and the Rost invariant of G takes values in $H^3(k, \mathbb{Z}/4) \oplus H^3(k, \mathbb{Z}/3) \oplus H^3(k, \mu_5^{\otimes 2})$, i.e., consists of a mod-4, mod-3, and mod-5 component. If the group G splits by a field extension of degree coprime to l , then the l -component of its Rost invariant is zero ($l \in \mathbb{Z}$).

8.6 Lemma. *Let G be a group of type E_8 and X the variety of Borel subgroups of G . If the even component of the Rost invariant of G is trivial, then the motive of X with $\mathbb{Z}/2$ - (or $\mathbb{Z}_{(2)}$ -)coefficients is a direct sum of twisted copies of a motive R whose Poincaré polynomial over any splitting field of G equals $1 + t^{15} \in \mathbb{Z}[t]$.*

Proof. Since we are talking about the even component of the Rost invariant and about the motives with $\mathbb{Z}/2$ -coefficients we may assume that the base field k does not have any proper finite field extension of odd degree.

Let X_i be the projective G -homogeneous variety of maximal parabolic subgroups of G of type i . If the Rost invariant of G is trivial, then so is the Rost invariant of $G_{k(X_i)}$. On the other hand, since the semisimple anisotropic kernel of $G_{k(X_i)}$ has rank at most 7, the group $G_{k(X_i)}$ must be split (see e.g. [Ga01, Theorem 0.5]). Thus, X_i is a generically split variety for all $i = 1, \dots, 8$.

The classification of generically split varieties of type E_8 (see [PS10, Theorem 5.7(7)]) and the classification of the J -invariants for groups of type E_8 (see [PSZ, §6, Table] or Example 8.3) immediately imply that the J -invariant of G modulo 2 equals either $J_2(G) = (0, 0, 0, 1)$ or $(0, 0, 0, 0)$.

In the first case we are done by formula (5). In the second case the variety of Borel subgroups is a direct sum of twisted Tate motives and the present lemma is trivial. $\square$

8.7 Corollary. *Let k be a field with $\text{char } k = 0$. Let G be a group of type E_8 over k with the trivial even component of the Rost invariant. Then there exists a functorial invariant $u \in H^5(k, \mathbb{Z}/2)$ of G such that for any field extension K/k the invariant $u_K = 0$ iff G splits over a field extension of K of odd degree.*

Proof. Let X be the variety of Borel subgroups of G . It is generically split, since over $k(X)$ the group G splits. By Lemma 8.6 the motive of X contains a binary direct summand R which supports zero-cycles.

Consider the image $\overline{\text{Ch}}(X)$ of the natural map $\text{Ch}(X) \rightarrow \text{Ch}(X_{k(X)})$. By [PSZ, Proposition 6.4] and [KM06, Theorem 5.8]

$$\min\{i \mid \overline{\text{Ch}}_i(X) \neq 0\} = 15.$$

By Corollary 7.2 there exists a smooth projective variety $\tilde{Y}$ of dimension 15 birational to some closed subvariety Y of X such that $\tilde{Y}_{k(X)}$ has a zero-cycle of odd degree. The converse obviously holds, i.e., $X_{k(\tilde{Y})}$ has a zero-cycle of odd degree.

Therefore by Lemma 7.4 the variety $\tilde{Y}$ has the same direct summand R . We are done by Theorem 6.1. $\square$

We come now to Serre's problem. Let G be a group of type E_8 over $\mathbb{Q}$ such that $G_{\mathbb{R}}$ is a compact Lie group. Let now $K/\mathbb{Q}$ be a field extension and $q = \langle\langle -1, -1, -1, -1, -1 \rangle\rangle$ a 5-fold Pfister form. J.-P. Serre posed in a letter to M. Rost written on June 23, 1999 the following problem:

Is it true that G_K is split if and only if q_K is hyperbolic?

M. Rost replied on July 2, 1999 proving that if q_K is hyperbolic, then G_K is split (see [GS09] for the proof). In what follows we give a positive answer to Serre's question.

8.8 Theorem. *Serre's question has a positive answer.*

Proof. The compact real group of type E_8 has Rost invariant zero (see [Ga09, 13.4]), so we may speak of its invariant $u \in H^5(\mathbb{R}, \mathbb{Z}/2) = \{0, (-1)^5\}$ constructed in Corollary 8.7.

If $u = 0$, then by Corollary 8.7 this compact group splits over a field extension of odd degree. Since $\mathbb{R}$ has only one field extension of odd degree, we come to a contradiction. Thus, $u = (-1)^5$.

Let now G be a group of type E_8 defined over $\mathbb{Q}$ that becomes compact E_8 over $\mathbb{R}$. The Rost invariant of G lies in

$$H^3(\mathbb{Q}, \mathbb{Z}/4) \oplus H^3(\mathbb{Q}, \mathbb{Z}/3) \oplus H^3(\mathbb{Q}, \mu_5^{\otimes 2}) = H^3(\mathbb{Q}, \mathbb{Z}/4),$$

and it follows from [Ar75, Satz 2] that $H^3(\mathbb{Q}, \mathbb{Z}/4)$ injects into $H^3(\mathbb{R}, \mathbb{Z}/4) = H^3(\mathbb{R}, \mathbb{Z}/2)$. Therefore, G has Rost invariant zero over $\mathbb{Q}$, and again we may speak of the invariant $u \in H^5(\mathbb{Q}, \mathbb{Z}/2)$ of the group G . But restriction identifies $H^5(\mathbb{Q}, \mathbb{Z}/2)$ with $H^5(\mathbb{R}, \mathbb{Z}/2)$ (see [Ar75, Satz 2]). So $u = (-1)^5$.

Finally, if G_K splits for some field extension $K/\mathbb{Q}$, then $u_K = 0$ by Corollary 8.7. We are done. $\square$

References

- [Ar75] J.Kr. Arason. *Primideale im graduierten Witttring und im mod 2 Cohomologiering*. Math. Z. **145** (1975), 139–143.
- [Br03a] P. Brosnan. *Steenrod operations in the Chow theory*. Trans. Amer. Math. Soc. **355** (2003), no. 5, 1869–1903.
- [Br03b] P. Brosnan. *A short proof of Rost nilpotence via refined correspondences*. Doc. Math. **8** (2003), 69–78.
- [Ga01] S. Garibaldi. *The Rost invariant has trivial kernel for quasi-split groups of low rank*. Comment. Math. Helv. **76** (2001), no. 4, 684–711.

- [Ga09] S. Garibaldi. *Orthogonal involutions on algebras of degree 16 and the Killing form of E_8* . Appendix by K. Zainoulline. In *Quadratic forms — algebra, arithmetic, and geometry* (Baeza, Chan, Hoffmann, Schulze-Pillot, eds.), Contemp. Math. **493** (2009), 131–162.
- [GS09] S. Garibaldi, N. Semenov. *Degree 5 invariant of E_8* . Preprint 2009, to appear in IMRN. Available from <http://arxiv.org/abs/0903.0304>
- [GMS] S. Garibaldi, A. Merkurjev, J.-P. Serre. *Cohomological invariants in Galois cohomology*. University Lecture Series 28, AMS, Providence, RI, 2003.
- [Inv] M.-A. Knus, A. Merkurjev, J.-P. Tignol, M. Rost. *The book of involutions*. AMS Colloquium Publications (1998).
- [IzhV00] O. Izhboldin, A. Vishik. *Quadratic forms with absolutely maximal splitting*. Contemp. Math. **272** (2000), 103–125.
- [KM06] N. Karpenko, A. Merkurjev. *Canonical p -dimension of algebraic groups*. Adv. Math. **205** (2006), no. 2, 410–433.
- [LM] M. Levine, F. Morel. *Algebraic cobordism*. Springer Verlag, 2007.
- [PS10] V. Petrov, N. Semenov. *Generically split projective homogeneous varieties*. Duke Math. J. **152** (2010), 155–173.
- [PSZ] V. Petrov, N. Semenov, K. Zainoulline. *J -invariant of linear algebraic groups*. Ann. Sci. Éc. Norm. Sup. **41** (2008), 1023–1053.
- [Ro98] M. Rost. *The motive of a Pfister form*. Preprint, 1998. Available from <http://www.math.uni-bielefeld.de/~rost>
- [Ro02] M. Rost. *Norm varieties and algebraic cobordism*. Proc. of the Int. Congress of Math., vol. 2 (Beijing, 2002), 77–85, Higher Ed. Press, Beijing, 2002.
- [Ro07] M. Rost. *On the basic correspondence of a splitting variety*. Preprint 2007. Available from <http://www.math.uni-bielefeld.de/~rost>

- [S08] N. Semenov. *Motivic decomposition of a compactification of a Merkurjev-Suslin variety*. J. reine und angew. Math. **617** (2008), 153–167.
- [Vi98] A. Vishik. *Integral motives of quadrics*. Max-Planck-Institut für Mathematik, 1998. Available from <http://www.mpim-bonn.mpg.de>
- [Vi07] A. Vishik. *Symmetric operations in algebraic cobordism*. Adv. Math. **213** (2007), 489–552.
- [ViZ08] A. Vishik, K. Zainoulline. *Motivic splitting lemma*. Doc. Math. **13** (2008), 81–96.
- [Vo95a] V. Voevodsky. *Triangulated category of motives over a field*. 1995. Available from <http://www.math.uiuc.edu/K-theory/0074>
- [Vo95b] V. Voevodsky. *Bloch-Kato conjecture for $\mathbb{Z}/2$ -coefficients and algebraic Morava K -theories*. 1995. Available from <http://www.math.uiuc.edu/K-theory/0076>
- [Vo96] V. Voevodsky. *Milnor conjecture*. 1996. Available from <http://www.math.uiuc.edu/K-theory/0170>
- [Vo01] V. Voevodsky. *On 2-torsion in motivic cohomology*. 2001. Available from <http://www.math.uiuc.edu/K-theory/0502>
- [Vo02] V. Voevodsky. *Cancellation theorem*. 2002. Available from <http://www.math.uiuc.edu/K-theory/0541>
- [Vo03] V. Voevodsky. *On motivic cohomology with $\mathbb{Z}/l$ -coefficients*. 2003. Available from <http://www.math.uiuc.edu/K-theory/0639>
- [Za09] K. Zainoulline. *Special correspondences and Chow traces of Landweber-Novikov operations*. J. für die reine und angew. Math. **628** (2009), 195–204.

NIKITA SEMENOV

JOHANNES GUTENBERG-UNIVERSITÄT MAINZ, INSTITUT FÜR MATHEMATIK, STAUDINGERWEG 9, D-55099 MAINZ, GERMANY
 semenov@uni-mainz.de